

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 47/12 // (A01N 47/12, 43:50, 25:22)		A1	(11) International Publication Number: WO 99/65316
			(43) International Publication Date: 23 December 1999 (23.12.99)
(21) International Application Number: PCT/US99/14171		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 18 June 1999 (18.06.99)			
(30) Priority Data: 09/099,726 19 June 1998 (19.06.98) US			
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/099,726 (CON) Filed on 19 June 1998 (19.06.98)			
(71) Applicant (for all designated States except US): LONZA INC. [US/US]; 17-17 Route 208, Fair Lawn, NJ 07410 (US)			
(72) Inventors; and (75) Inventors/Applicants (for US only): LUTZ, Patrick, J. [US/US]; 5735 Kesslerville Road, Nazareth, PA 18064 (US). BAN, Susan, Alcorn [US/US]; Rural Road #3, P.O. Box 36A, Church Lane, Kunkletown, PA 18058 (US). FARINA, Thomas, E. [US/US]; 14 Glenn Road, Flemington, NJ 08822 (US)		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(74) Agents: LEWEN, Bert, J. et al.; Darby & Darby P.C., 805 Third Avenue, New York, NY 10022-7513 (US).			
(54) Title: STABILIZED MIXTURES OF AN IODOPROPARGYL COMPOUND AND A FORMALDEHYDE DONOR			
(57) Abstract <p>A highly stable liquid formulation having broad spectrum preservative properties which constitutes an admixture of a dialkanol-substituted DMH, an iodopropynyl compound, a stabilizer of a hydantoin, urea or derivative thereof, and a hydroxyl solvent. Preferably the constituents are dimethyloldimethylhydantoin, 3-iodo-2-propynylbutyl carbamate, dimethylhydantoin, and a glycol solvent. The preservative preferably has a total formaldehyde content of 5 % and less than 0.2 % of free formaldehyde. The composition is prepared by successively admixing the dialkanol-substituted dimethylhydantoin and the stabilizer, the hydroxyl solvent, and the iodopropynyl compound. Also described is a stabilized iodopropynyl compound preferably containing dimethylhydantoin as the stabilizer.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

STABILIZED MIXTURES OF AN IODOPROPARGYL COMPOUND AND A FORMALDEHYDE DONOR

Background of the Invention

10 The need for effective and economical preservative compositions is well known. There are a wide variety of applications where inhibiting the growth of microorganisms is necessary, as for example personal care products such as shampoos, creams, lotions, cosmetics, soap and household products such as laundry detergents, hard surface cleaners, and fabric softeners. The shelf life of these
15 preparations depends on their resistance to microbial spoilage.

 In addition, in many industrial applications, antimicrobial agents are useful in paint, wood, textiles, adhesives, sealants, leather, rope, paper pulp, plastics, fuel, oil, and rubber and metal working fluids. The control of slime-producing bacteria and fungi in pulp and paper mills and cooling towers is a matter
20 of substantial commercial importance.

 For the foregoing applications the demand for stable broad-spectrum preservatives has increased. In recent years, these needs have been met with solid formulations. For example, combinations of formaldehyde donors (e.g., dimethyloldimethylhydantoin (DMDMH)) and halopropynyl compounds (e.g., 3-
25 iodo-2-propynylbutyl carbamate (IPBC)) have achieved considerable commercial success. Such synergistic combinations have been described in U.S. Patent 4,844,891.

 Furthermore, because of the demand of governmental regulations, low free-formaldehyde products are needed. Research in this area has also proved
30 beneficial. For example, in the case of DMDMH, improved formulation and

processing has resulted in compositions which contain less than 0.1% free formaldehyde. (See U.S. Patent 5,405,862.) In contrast, earlier formulations of DMDMH had over 1% of free formaldehyde. (See U.S. Patent 3,987,184.)

At the present time, in addition to meeting the above criteria, the industry is demanding liquid forms of preservatives as the use of automatic liquid blending systems becomes more popular. Unfortunately, preservatives that are in liquid form, highly stable, broad spectrum, and low in free formaldehyde have eluded formulators.

10 Summary of the Invention

It has now been discovered that highly stable, liquid formulations of broad spectrum preservatives can be prepared by admixing alkanol-substituted dimethylhydantoins, iodopropynyl compounds, stabilizers, and a solvent. This invention is based, in part, on the unexpected finding that the iodopropynyl compounds can be stabilized at high temperatures by the addition of hydantoin-type stabilizers such as dimethylhydantoin. Furthermore, the compositions of the invention have surprisingly good physical stability at low temperatures and enhanced solubility which allows the easy preparation of these highly concentrated mixtures. Such highly concentrated preservatives useful in automatic liquid blending systems could not heretofore be prepared.

Brief Description of the Figures

Figure 1 illustrates the results of a 48 hour minimum inhibitory concentration test against 8 bacteria organisms of the liquid formulation of the invention and a solid formulation of DMDMH and IPBC.

Figure 2 illustrates the results of a 72 hour minimum inhibitory concentration test against 4 fungal organisms of the liquid formulation of the invention and a solid formulation of DMDMH and IPBC.

Detailed Description of the Invention


	Broad wt. %	Preferred wt. %
Dialkanol-substituted DMH	20-95 %	75-85 %
Iodopropynyl Compound	0.2-20	1-5
Stabilizer	1-30	5-20
Hydroxyl Solvent	0-60	2-20

The ratio of the stabilizer to the iodopropynyl compound may broadly be from about 150:1 to 0.05:1, preferably from 20:1 to 1:1, most desirably from about 10:1 to 2:1.

The alkanol-substituted DMH compounds used in the invention are well known and include those defined in U.S. Patent Nos. 3,987,184 and 4,172,140, the entire contents of which are incorporated herein by reference. These are condensation products of 5,5-dimethylhydantoin with 1, 2, or more moles of formaldehyde (e.g., 1,3-dimethylol-5,5-dimethylhydantoin, 1-methylol-5,5-dimethylhydantoin, or 3-methylol-5,5-dimethylhydantoin, 1,3-dimethylol-oxymethylene-5,5-dimethylhydantoin, 1-methylol-3-methyloloxymethylene-5,5-dimethylhydantoin and 1,3-dimethyloloxymethylene-5,5-dimethylhydantoin, and mixtures thereof).

Examples of compounds which may be used as the iodopropynyl component of the invention are the fungicidally active iodopropynyl derivatives. These include compounds derived from propynyl or iodopropynyl alcohols such as the esters, ethers, acetals, carbamates and carbonates and the iodopropynyl derivatives of pyrimidines, triazolinones, tetrazoles, triazinones, sulfamides, benzothiazoles, ammonium salts, carboxamides, hydroxamates, and ureas. Preferred among these compounds is 3-iodo-2-propynylbutyl carbamate, IPBC. These compounds are included within the broadly useful class of compounds having the generic formulas such as I and II shown below:

4




wherein:

R is selected from the group consisting of substituted and
 10 unsubstituted alkyl, aryl, and alkylaryl groups having from 1 to 20 carbon atoms;
 and

m and n are independent integers from 1 to 3.

15



20

wherein:

R₁ and R₂ are defined as R₃ and R₄ below or are joined to form a
 cycloalkyl, cycloalkenyl, aromatic or a heterocyclic ring containing an oxygen,
 25 nitrogen or sulfur atom or an alkoxy, amino, carbonyl, carboxyl, halo, hydroxyl,
 keto or a thiocarboxyl-substituted derivative thereof;


R₃, R₄ and R₅ are independently selected from (A) hydrogen, alkyl,
 cycloalkyl, alkenyl, cycloalkenyl, aryl, a heterocyclic ring containing an oxygen,
 nitrogen or sulfur atom, alkoxy, amino, carbonyl, carboxyl, halo, hydroxyl, keto or
 30 a thiocarboxyl and (B) substituted derivatives of the alkyl, cycloalkyl, alkenyl,
 cycloalkenyl, aryl and the heterocyclic ring wherein the substitutions are alkyl,
 cycloalkyl, alkenyl, cycloalkenyl, aryl, alkoxy, amino, carboxyl, halo, hydroxyl,
 keto or a thiocarboxyl;

a is 0 to 16;

W may be a single bond, oxygen, NR_6 , or $(\text{CR}_7\text{R}_8)_m$, wherein R_6 is hydrogen, alkyl, cycloalkyl, alkenyl, cycloalkenyl, aryl or a heterocyclic ring containing an oxygen, nitrogen or sulfur atom or a substituted derivative of alkyl, cycloalkyl, alkenyl, cycloalkenyl or aryl groups wherein the substitutions are alkyl, cycloalkyl, alkenyl, cycloalkenyl, aryl, alkoxy, amino, carbonyl, carboxyl, halo, hydroxyl, keto, or a thiocarboxyl wherein R_6 , R_7 and R_8 are defined as R_3 and R_4 above and m is an integer from 1 to 12. The above definition of R_6 includes, among other things, an aminoalkyl group.

The heterocyclic rings referred to in the above definitions may contain from 5 to 8 members, the alkyl or cycloalkyl groups from 1 to 18 atoms, the alkenyl or cycloalkenyl groups from 2 to 18 carbon atoms, and the aryl groups from 6 to 10 members.

In formula II, when R_1 and R_2 are hydrogen, R_3 and R_4 are carbonyl, R_5 is $-\text{CH}=\text{CH}-\text{CO}_2\text{H}$; a is equal to 0; and W is oxygen, the compound is iodopropynyl maleate,




Other compounds include the mono-iodopropynyl esters of anhydrides such as succinic and phthalic as well as the following anhydrides: ethylenediamine tetraacetic dianhydride, 3,3-dimethylglutaric anhydride, S-acetylmercaptosuccinic anhydride, dichloromaleic anhydride, 2-dodecen-1-yl succinic anhydride and cis-5-norbornene-endo-2,3-dicarboxylic anhydride. Where hydrophilicity is desired, the sodium salts may be used because of their extremely high water solubility. Preferred carboxylic acid anhydrides include succinic, itaconic, phthalic, tetrachlorophthalic, and diglycolic anhydride. Such compounds are defined in U.S. Patent 4,844,891 and 5,073,570.

The stabilizers used in the invention are hydantoin and urea and their derivatives, most preferably 5,5-dimethylhydantoin.


6

Hydantoins and their
derivatives may be
represented by formu-
las III, IV, and V:



III

10



IV

where R_1 to R_4 are independently selected from H, C_1 to C_{22} .

N,N''-Methylenebis[N'-2,5-dioxo-4-imidazolidinyl]urea and its
derivatives:

15


where R_1 to R_5 are independently selected from H or C_1 to C_{22} .

2,5-Dioxo-4-imidazolidinyl urea (5-ureidohydantoin) and its deriva-
tives:

7

where R₁ to R₇ are independently selected from H, CH₃, C₂H₅ or C₃H₇.

Urea and its derivatives are represented by Formula VI:




where R_1 to R_4 are independently selected from H or C_1 to C_{12} . Where all the R groups are H, the compound is urea.

The solvents which may be used in the invention include mono-, di-, and polyhydroxyl alcohols. For example, monohydroxyl alcohols having from about 1 to 5 carbon atoms, most preferably ethanol and propanol, may be used. Dihydroxyl alcohols (i.e., glycols) such as C₂ to C₈ diols such as propylene glycol and butylene glycol are advantageous. 1,3-Butylene glycol is particularly preferred. Other compounds which can be used include: dipropylene glycol, glycerin, diglycerin, PPG-9, PPG-2-buteth-2, butoxypropanol, butoxydiglycol, PPG-2 butyl ether, glycereth-7, sorbitol, isopentyldiol, myristyl myristate, and phenoxy ethanol.

The preservative formulations of the instant invention can be readily prepared in ac-

25 cordance with
procedures well
known to those
skilled in the
art. The pre-



ferred procedure is first to mix the stabilizer at temperatures ranging from 30°C to 50°C with the dialkanol-substituted dimethylhydantoin. This mixture is stirred for 5 minutes at 30°C. It may be heated to 50°C to increase the solution rate.

Thereafter the hydroxylic solvent is added and the entire mixture is stirred over a
5 period of 5 minutes. Finally, the iodopropynyl compound is added and mixed for another 15 minutes to form a homogeneous solution. The total mixing time is approximately 30 minutes.

The liquid preservative composition of the invention has a free formaldehyde concentration of less than 1 wt.%, preferably less than 0.2. The total
10 formaldehyde concentration is from 5 wt.% to 25 wt.%, and preferably from 12 wt.% to 14 wt.%.

The preservatives of the invention can be used as active compounds for combating microorganisms, in particular for the preservation of cosmetics, personal care products, household products, and industrial materials such as
15 adhesives, sizes, paper and cardboard, textiles, leather, wood, paints and articles made of plastic, cooling lubricants and other materials which can be attacked or decomposed by microorganisms. Components of production plants, for example cooling water, which can be impaired by multiplication of microorganisms, may also be beneficially treated. Also, the integrity of other water-containing systems,
20 such as swimming pools and spas, can be maintained by use of the preservatives of the invention. In addition, they can be used to control and eliminate microorganisms by disinfection and sanitization of surfaces, such as found in homes, institutions, and hospitals.

Examples of microorganisms which can effect contamination,
25 degradation, or a change in the industrial environments and materials are bacteria, fungi, yeasts, algae, and slime organisms. The active compounds of the invention act against fungi, in particular mold fungi, fungi which discolor and destroy wood (Basidiomycetes), and against slime organisms and algae.

Microorganisms of the following genera are examples: *Alternaria*, such as *Alternaria tenuis*, *Aspergillus*, such as *Aspergillus niger*, *Chaetomium*, such as *Chaetomium globosum*, *Candida*, such as *Candida albicans*, *Lentinus*, such as *Lentinus tigrinus*, *Penicillium*, such as *Penicillium glaucum*, *Trichophyton*, such as *Trichophyton mentagrophytes*, *Aureobasidium*, such as *Aureobasidium pullulans*, *Enterobacter*, such as *Enterobacter gergoviae*, *Trichoderma*, such as *Trichoderma viride*, *Escherichia*, such as *Escherichia coli*, *Pseudomonas*, such as *Pseudomonas aeruginosa* and *Pseudomonas cepacia*, and *Staphylococcus*, such as *Staphylococcus aureus* and *Staphylococcus epidermidis*.

The use concentrations of the active compounds according to the invention depend on the nature and the occurrence of the microorganisms to be combated, and on the composition of the material to be preserved. The optimum amount to be employed can be determined by means of a series of tests. The use concentrations are in general in the range of 0.00005 (0.5 ppm) to 5% by weight, preferably from 0.0001 to 1%, relative to the material to be preserved.

Liquid compositions of this invention are used directly as they are manufactured without dilution. They may be poured into small batches (from one to thousands of pounds) of product at any point in its manufacture. Also, the liquid compositions may be pumped into medium sized batches (from thousands to tens of thousands of pounds) from a weigh scale.

The preservative of the invention may also be metered continuously from a storage tank into large sized production runs (from tens of thousands to millions of pounds) in systems custom-designed to continuously mix all the components of the finished product at approximately the same rate that it is filled into its final package. The blending elements of continuous mixers are mostly shaped in the form of spirals or screws, effecting on rotation both a mixing and a transport of the product composition.

Because start-up is very labor-intensive, to insure all the metering equipment is properly calibrated, these systems are generally used only for very high volume, long and continuous production runs.

In order to illustrate more fully the subject invention, attention is
5 directed to the following examples:

Example 1

A preservative of the instant invention containing 81 parts of Glydant II (a trademark of Lonza Inc.), 12 parts of dimethylhydantoin, 4.5 parts of butylene glycol, and 2.5 parts of iodopropynyl butyl carbamate is described in this example.
10 Glydant II contains 65% DMDMH, 30% MMDMH, and 5% DMH. It has a total formaldehyde content of 17%.

Initially the DMH and the Glydant II are mixed at 30°C. The mixing continues for 5 minutes during which the temperature is increased to 50°C.
15 Thereafter the butylene glycol is added and stirring is continued for 10 minutes. Finally, the IPBC is added and mixed with the other constituents for 15 minutes to form the final solution. After the mixture is cooled to room temperature, a clear homogeneous liquid solution is obtained. The solution has a total formaldehyde content of 14 wt. %, a free formaldehyde content of 0.05 wt. %, and an IPBC
20 content of 2.3%. After storage for one month at 50°C, the mixture remains a clear, colorless liquid with 92% recovery of the total formaldehyde content and > 99% recovery of the IPBC content obtained.

This solution remains clear and colorless even after two months storage at room temperature in sunlight. Significantly, quantitative recovery of total
25 formaldehyde and IPBC is also obtained. Free formaldehyde content is 0.06%.

Example 2

To illustrate the suitability of the liquid preservative of the invention for use in automated liquid blending systems, the viscosity and the specific gravity

of the formulation described in Example 1 are determined. A Brookfield Model RVT viscometer, Spindle No. 3, at 50 rpm is used to determine viscosity. The following results are obtained:

Table 1

<u>Temperature</u>	<u>Specific Gravity</u>	<u>Viscosity, cps</u>
5°C		120
15°C	1.2043	-
25°C	1.1992	40
30°C	1.1960	-
35°C	1.1954	-
45°C		30
65°C	1.1625	20
85°C	1.1500	10

The viscosity values and the specific gravity measurements presented in Table 1 for the liquid preservative of this invention are typical of liquids used in automated blending systems. These ideal parameters are dependent on the particular system employed.

Example 3

To demonstrate the activity of the liquid preservative described in Example 1, 48 hour minimum inhibitory concentration tests were performed on eight bacteria organisms. These tests compared the liquid preservative of the invention to a solid preservative on a 100% active basis. The composition of the solid preservative was 95% DMDMH and 5% IPBC.

The results obtained from the foregoing tests are set forth in Figure

1.

It will be noted that the results obtained for the liquid formulation of the invention, as compared to the solid formulation, are substantially comparable. In certain instances the liquid formulation is superior, as in the case of the *A. baumannii*, the *S. epidermidis*, and the *E. coli*. These data show that the liquid formulations of the invention have a broad spectrum activity against a wide range of bacteria.

Example 4

Minimum inhibitory concentration tests (72 hours) for the liquid preservative described in Example 1 and the solid preservative defined in Example 2 were performed with respect to four fungal organisms. The comparison was made on a 100% active basis.

Figure 2 illustrates that the liquid preservative of the invention is as efficacious as the solid in all of the tests performed and that these materials have broad spectrum activity against a variety of fungi.

Example 5

A series of formulations were prepared to determine the solubility, free formaldehyde concentration, and physical stability of the compositions of the invention as compared to compositions free of dimethylhydantoin. The formulations are shown in Table 2.

Table 2
Liquid Glydant Plus Comparison Chart

Ingredients - %	A 5503-84-1	B 5503-84-2	C 5503-90-1	D 5503-111-1	E 5503-111-2	F 5503-115-1	G 5503-115-2
IPBC (Poly P-100)	2.50	2.50	2.50	2.50	2.50	2.50	2.50
Glydant II	74.70	72.85	74.70	81.00	81.00	81.00	81.00
Prop. Glycol	22.80		12.80	4.50		16.50	4.50
But. Glycol		24.65			4.50		
DMH			10.00	12.00	12.00		12.00
DI Water							No good-ppt.
Solubility	OK	OK	OK	OK	OK		
Physical Solubility	ppt. after 3rd freeze/thaw	OK after 3rd freeze/thaw	OK after 3rd freeze/thaw	OK after 3rd freeze/thaw	OK after 3rd freeze/thaw	ppt. after 3rd freeze/thaw	
% Free HCHO - Initial	0.20	0.24	0.07	0.03	0.05	0.17	
% Free HCHO - 1 month @ 40°C	0.19	0.19	0.19	0.02	0.03		
% Free HCHO - 2 months @ 40°C	0.21	0.25	0.06				
% Free HCHO - 1 month @ 50°C	0.24	0.23	0.11	0.05	0.05		
Comments	Unstable & % free HCHO too high	Stable but % free HCHO too high	Stable & good % free HCHO but % total HCHO not high enough	Stable & low % free HCHO & % total HCHO - OK	Stable & low % free HCHO & total HCHO - OK Going with But. Glycol	Unstable and % free HCHO too high	Insoluble

5

10

15

20

The above data show that the liquid compositions of the invention, namely, Formulations 90-1, 111-1, and 111-2, had high solubility, good physical stability, and extremely low free formaldehyde as compared to the formulations free of DMH, namely, Formulations 84-1, 84-2, 115-1, and 115-2. Formulations 111-1 and 111-2 are preferred because of the higher total formaldehyde concentration.

Example 5

Using the formulations shown in Table 3, liquid preservatives of the invention were tested to determine the free formaldehyde, total formaldehyde, and percent IPBC after one month stability tests at 40°C and 50°C, respectively. As will be noted, the formulations contained 10, 12, and 15 parts of dimethylhydantoin. Results are set forth in the following table:

Table 3

Chemical Stability Results of the Preferred Aqueous Liquid Systems of the Invention

Formulation	A 5503- 109-1	B 5503- 109-2	C 5503- 111-1	D 5503- 111-2	E 5503- 111-3
Parts:					
Glydant II	81	81	81	81	81
IPBC	2.5	2.5	2.5	2.5	2.5
Propylene glycol	6.5	-	4.5	-	1.5
Butylene glycol	-	6.5	-	4.5	-
DMH	10	10	12	12	15
<u>% Free HCHO</u> (criterion for Free HCHO is <0.19%)					
Initial	0.05	0.07	0.03	0.05	0.02
40°C Stability 1 Month	0.05	0.05	0.02	0.03	0.03
50°C Stability 1 month	0.07	0.05	0.05	0.05	0.02
<u>% Total HCHO</u> (criterion for Total HCHO is 90% recovery)					

	Initial	14.08	14.05	14.07	14.09	14.11
	40°C Stability 1 Month	13.04	13.10	12.33	13.16	13.19
	50°C Stability 1 Month	12.83	13.07	12.80	12.91	12.84
5	% Recovery after 50°C Stability	91.0	93.0	91.0	92.0	91.0
	% IPBC (criterion for IPBC is 90% recovery)					
	Initial	2.41	2.32	2.34	2.28	2.46
	40°C Stability 1 Month	2.38	2.38	2.48	2.42	2.43
10	50°C Stability 1 Month	2.23	2.20	2.24	2.27	2.26
	% Recovery after 50°C Stability	93.0	95.0	96.0	99.6	92.0

15 The above data show that, after one month of storage, the stabilized compositions all had stabilities of over 90%. These data illustrate that butylene glycol is the preferred solvent.

 In comparison to a liquid preservative system, without the added stabilizer, DMH, recovery of IPBC after four weeks storage at 45°C was only about
20 60% and does not meet the criteria of the industries that use preservatives in their products.

Example 6

25 The composition of the invention described in Example 1 was tested for chemical stability at elevated temperatures and in sunlight in glass and high density polyethylene containers. The compositions were analyzed for percent total formaldehyde, percent free formaldehyde, and percent IPBC after one, two, and three months.

The results are shown in the following table:

Table 4

		1 month		2 months		3 months	
		glass	HDPE	glass	HDPE	glass	HDPE
5	#5503-125	Initial					
	% Total HCHO	13.02					
	RT		13.60	13.40	13.90	12.80	13.11
	40°C		13.40	13.60	13.10	13.10	12.76
	50°C		13.10	13.10	--	--	--
10	Sunlight		13.60	13.60	13.50	13.20	12.95
	Lab light		--	--	--	--	12.93
	% Free HCHO	0.09					
	RT		0.05	0.04	0.04	0.04	0.02
	40°C		0.04	0.06	0.06	0.05	0.04
15	50°C		0.06	0.05	--	--	--
	Sunlight		0.05	0.06	0.09	0.06	0.07
	Lab light		--	--	--	--	0.03
	% IPBC	2.49					
	RT		2.50	2.50	2.30	2.30	2.30
20	40°C		2.49	2.45	2.50	2.50	2.40
	50°C		2.26	2.25	--	--	--
	Sunlight		2.46	2.49	2.30	2.50	2.30
	Lab light		--	--	--	--	2.30

25

The above data clearly show that the composition remains stable under all conditions, i.e., at room temperature, 40°C, and 50°C and in the presence of sunlight and laboratory lighting.

We claim:

- 1 1. A broad spectrum liquid preservative formulation comprising from
2 about 20 to 95 parts of a dialkanol-substituted dimethylhydantoin, from about 0.2 to
3 20 parts of an iodopropynyl compound, from about 1 to 30 parts of a stabilizer for the
4 iodopropynyl compound, and from about 0 to 60 parts of a hydroxyl solvent.
- 1 2. The liquid preservative formulation of claim 1 wherein the dialkanol-
2 substituted dimethylhydantoin is 1,3-dimethylol-5,5-dimethylhydantoin, 1-methylol-
3 5,5-dimethylhydantoin, 3-methylol-5,5-dimethylhydantoin, 1,3-dimethylol-
4 oxymethylene-5,5-dimethylhydantoin, 1-methylol-3-methylloxymethylene-5,5-
5 dimethylhydantoin and 1,3-dimethylloxymethylene-5,5-dimethylhydantoin, and
6 mixtures thereof.
- 1 3. The liquid preservative formulation of claim 1 wherein the dialkanol
2 disubstituted dimethylhydantoin is 1,3-dimethylol-5,5-dimethylhydantoin.
- 1 4. The liquid preservative formulation of claim 1 wherein the
2 iodopropynyl compound is 3-iodo-2-propynylbutyl carbamate.
- 1 5. The liquid preservative formulation of claim 1 wherein the stabilizer is
2 hydantoin, urea, or a derivative thereof.
- 1 6. The liquid preservative formulation of claim 5 wherein the stabilizer is
2 dimethylhydantoin.
- 1 7. The liquid preservative formulation of claim 1 wherein the hydroxyl
2 solvent is propylene glycol or butylene glycol.
- 1 8. The liquid preservative formulation of claim 6 wherein the ratio of the
2 dimethylhydantoin to the iodopropynyl compound is from 150:1 to 0.05:1.

1 9. The liquid preservative formulation of claim 1 wherein the free
2 formaldehyde concentration is less than 1 % and the total formaldehyde concentration
3 at least 2 %.

1 10. A method of preparing a liquid preservative composition which
2 comprises blending a dialkanol-substituted dimethylhydantoin and a stabilizer of
3 hydantoin, urea or a derivative thereof to form a homogeneous mixture, admixing a
4 hydroxyl solvent and the iodopropynyl compound with the foregoing mixture to
5 obtain a homogeneous solution containing a total formaldehyde content of at least 2 %
6 and less than 0.2 % free formaldehyde.

1 11. A stabilized iodopropynyl solution which comprises an iodopropynyl
2 compound and a stabilizing amount of a hydantoin, urea or derivative thereof.

1 12. The solution of claim 11 wherein the stabilizer is dimethylhydantoin.

1 13. The solution of claim 11 wherein the iodopropynyl compound is iodo-
2 2-propynylbutyl carbamate.

1 14. A process for killing or retarding the growth of microbes which
2 comprises admixing the composition of claim 1 in a solution subject to the growth of
3 microbes.

1 15. The process of claim 14 wherein the composition contains a dialkanol-
2 substituted dimethylhydantoin, a stabilizer of hydantoin, urea or a derivative thereof, a
3 hydroxyl solvent, and an iodopropynyl compound.

2
AS FILED 5/19/98
Solid and Liquid Glycyl Plus (100% active basis) ppm

FIGURE 1
48 HOUR MIC RESULTS FOR LIQUID GLYDANT PLUS VS.
SOLID GLYDANT PLUS ON 100% ACTIVE BASIS AGAINST
8 BACTERIA ORGANISMS





FIGURE 2
 72 HOUR MIC RESULTS FOR LIQUID GLYDANT PLUS vs.
 SOLID GLYDANT PLUS ON 100% ACTIVE BASIS AGAINST
 4 FUNGAL ORGANISMS



INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/14171

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A01N47/12 //(A01N47/12,43:50,25:22)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Reference of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	DATABASE WPI Section Ch, Week 199834 Derwent Publications Ltd., London, GB; Class D22, AN 1998-393347 XP002118218 & JP 10 158110 A (MITSUI TOATSU CHEM INC), 16 June 1998 (1998-06-16) abstract & PATENT ABSTRACTS OF JAPAN vol. 199, no. 811, 30 September 1998 (1998-09-30) & JP 10 158110 A abstract	11,13
A	WO 95 29588 A (ISP CHEMICALS INC) 9 November 1995 (1995-11-09) page 3, paragraph 3 -page 5, paragraph 1 --- -/-	1-15

☒ Further documents are listed in the continuation of box C☒ Patent family members are listed in annex

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search

8 October 1999

Date of mailing of the international search report

18/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lamers, W

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/14171

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 327 220 A (LONZA AG) 9 August 1989 (1989-08-09) cited in the application page 3, line 20 - line 48 ---	1-15
A	WO 96 39836 A (TROY CHEMICAL COMPANY) 19 December 1996 (1996-12-19) page 1, line 5 -page 14, line 21 ---	1-15
A	EP 0 571 903 A (LONZA AG) 1 December 1993 (1993-12-01) cited in the application page 2, line 3 -page 3, line 5 -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/14171

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
JP 10158110	A	16-06-1998	NONE	
WO 9529588	A	09-11-1995	US 5428050 A	27-06-1995
			US 5552425 A	03-09-1996
			AU 691613 B	21-05-1998
			AU 2392395 A	29-11-1995
			CA 2183652 A	09-11-1995
			EP 0757518 A	12-02-1997
			HU 76978 A	28-01-1998
			NZ 284708 A	29-01-1997
			PL 316963 A	17-02-1997
			US 5496842 A	05-03-1996
			US 5631273 A	20-05-1997
EP 0327220	A	09-08-1989	US 4844891 A	04-07-1989
			AT 93681 T	15-09-1993
			AU 2958089 A	03-08-1989
			CA 1331141 A	02-08-1994
			CN 1036882 A, B	08-11-1989
			DE 68908730 D	07-10-1993
			DE 68908730 T	27-01-1994
			DK 47189 A	04-08-1989
			ES 2059715 T	16-11-1994
			JP 1940584 C	09-06-1995
			JP 2191204 A	27-07-1990
			JP 6067808 B	31-08-1994
			KR 9506926 B	26-06-1995
WO 9639836	A	19-12-1996	AU 5986196 A	30-12-1996
			CA 2223778 A	19-12-1996
			EP 0831706 A	01-04-1998
			NO 975675 A	06-02-1998
EP 0571903	A	01-12-1993	AT 141919 T	15-09-1996
			AU 3872693 A	02-12-1993
			BR 9302029 A	30-11-1993
			CA 2093045 A, C	22-11-1993
			CN 1087778 A	15-06-1994
			DE 69304253 D	02-10-1996
			DE 69304253 T	30-01-1997
			DK 571903 T	10-02-1997
			ES 2093878 T	01-01-1997
			GR 3021711 T	28-02-1997
			HU 64177 A	28-12-1993
			JP 2657148 B	24-09-1997
			JP 6179658 A	28-06-1994
			KR 9702875 B	12-03-1997
			PL 299028 A	10-01-1994
			PL 173873 B	29-05-1998
			US 5405862 A	11-04-1995
			ZA 9303464 A	20-12-1993

